

Glycemic Index by Glycemic Load

--	LOW GI	MED GI	HI GI
LOW GL	All-bran cereal (8,42) Apples (6,38) Carrots (3,47) Chana dal (3,8) Chick peas (8,28) Grapes (8,46) Kidney beans (7,28) Nopal (0,7) Oranges (5,42) Peaches (5,42) Peanuts (1,14) Pears (4,38) Pinto beans (10,39) Red lentils (5,26) Strawberries (1,40) Sweet corn (9,54)	Beets (5,64) Cantaloupe (4,65) Pineapple (7,59) Sucrose (table sugar) (7,68)	Popcorn (8,72) Watermelon (4,72) Whole wheat flour bread (9,71) White wheat flour bread (10,70)
MED GL	Apple juice (11,40) Bananas (12,52) Buckwheat (16,54) Fettucine (18,40) Navy beans (12,38) Orange juice (12,50) Parboiled rice (17,47) Pearled barley (11,25) Sourdough wheat bread (15,54)	Life cereal (16,66) New potatoes (12,57) Sweet potatoes (17,61) Wild rice (18,57)	Cheerios (15,74) Shredded wheat (15,75)
HI GL	Linguine (23,52) Macaroni (23,47) Spaghetti (20,42)	Couscous (23,65) White rice (23,64)	Bagel (25,72) Baked Russet potatoes (26,85) Cornflakes (21,81)

GI: low=1-55 mid=56-69 High=70-100 GL: low=1-10 mid=11-19 High=20 or more

This article originally appeared on [Mendosa.com](http://www.mendosa.com) on April 23, 2003.

copied from http://www.mendosa.com/common_foods_bw.htm